

Complex Trigonometric and Hyperbolic Functions (7A)

-
-


Copyright (c) 2011 - 2015 Young W. Lim.

Permission is granted to copy, distribute and/or modify this document under the terms of the GNU Free Documentation License, Version 1.2 or any later version published by the Free Software Foundation; with no Invariant Sections, no Front-Cover Texts, and no Back-Cover Texts. A copy of the license is included in the section entitled "GNU Free Documentation License".

Please send corrections (or suggestions) to youngwlim@hotmail.com.

This document was produced by using OpenOffice and Octave.

cosh(x)




$$\cosh \alpha = \frac{1}{2}(e^{+\alpha} + e^{-\alpha})$$

$$\cos \theta = \frac{1}{2}(e^{+j\theta} + e^{-j\theta})$$

<http://en.wikipedia.org/>

$\sinh(x)$




$$\sinh \alpha = \frac{1}{2}(e^{+\alpha} - e^{-\alpha})$$

$$\sin \theta = \frac{1}{2j}(e^{+j\theta} - e^{-j\theta})$$

<http://en.wikipedia.org/>

Definitions of Hyperbolic Functions



$$\cosh \alpha = \frac{1}{2}(e^\alpha + e^{-\alpha})$$

$$\sinh \alpha = \frac{1}{2}(e^\alpha - e^{-\alpha})$$

$$\tanh \alpha = \frac{(e^\alpha - e^{-\alpha})}{(e^\alpha + e^{-\alpha})}$$

<http://en.wikipedia.org/>

Hyperbolic vs. Trigonometric Functions

Trigonometric Function

$i x$

$$e^{+ix} = \cos x + i \sin x$$

$$e^{-ix} = \cos x - i \sin x$$

$$\cos x = \frac{1}{2}(e^{+ix} + e^{-ix})$$

$$\sin x = \frac{1}{2i}(e^{+ix} - e^{-ix})$$

$$\tan x = \frac{1}{i} \frac{(e^{+ix} - e^{-ix})}{(e^{+ix} + e^{-ix})}$$

Hyperbolic Function

x

$$e^{+x} = \cosh x + \sinh x$$

$$e^{-x} = \cosh x - \sinh x$$

$$\cosh x = \frac{1}{2}(e^{+x} + e^{-x})$$

$$\sinh x = \frac{1}{2}(e^{+x} - e^{-x})$$

$$\tanh x = \frac{(e^{+x} - e^{-x})}{(e^{+x} + e^{-x})}$$

Trigonometric functions with imaginary arguments

$$ix \rightarrow x$$

$$\cos x = \frac{1}{2}(e^{+ix} + e^{-ix})$$

$$\sin x = \frac{1}{2i}(e^{+ix} - e^{-ix})$$

$$\tan x = \frac{1}{i} \frac{(e^{+ix} - e^{-ix})}{(e^{+ix} + e^{-ix})}$$

$$\cosh x = \frac{1}{2}(e^{+x} + e^{-x})$$

$$\sinh x = \frac{1}{2}(e^{+x} - e^{-x})$$

$$\tanh x = \frac{(e^{+x} - e^{-x})}{(e^{+x} + e^{-x})}$$

$$\cos ix = \cosh x$$

$$\sin ix = i \sinh x$$

$$\tan ix = i \tanh x$$

$$\cos ix = \frac{1}{2}(e^{-x} + e^{+x})$$

$$\sin ix = \frac{1}{2i}(e^{-x} - e^{+x})$$

$$\tan ix = \frac{1}{i} \frac{(e^{-x} - e^{+x})}{(e^{-x} + e^{+x})}$$

$$\cosh x = \frac{1}{2}(e^{+x} + e^{-x})$$

$$\sinh x = \frac{1}{2}(e^{+x} - e^{-x})$$

$$\tanh x = \frac{(e^{+x} - e^{-x})}{(e^{+x} + e^{-x})}$$

Hyperbolic functions with imaginary arguments

$$\cos x = \frac{1}{2}(e^{+ix} + e^{-ix})$$

$$\sin x = \frac{1}{2i}(e^{+ix} - e^{-ix})$$

$$\tan x = \frac{1}{i} \frac{(e^{+ix} - e^{-ix})}{(e^{+ix} + e^{-ix})}$$



$$\cosh x = \frac{1}{2}(e^{+x} + e^{-x})$$



$$\sinh x = \frac{1}{2}(e^{+x} - e^{-x})$$



$$\tanh x = \frac{(e^{+x} - e^{-x})}{(e^{+x} + e^{-x})}$$

$x \leftarrow ix$

$$\cos x = \frac{1}{2}(e^{+ix} + e^{-ix})$$

$$\sin x = \frac{1}{2i}(e^{+ix} - e^{-ix})$$

$$\tan x = \frac{1}{i} \frac{(e^{+ix} - e^{-ix})}{(e^{+ix} + e^{-ix})}$$



$$\cosh ix = \frac{1}{2}(e^{+ix} + e^{-ix})$$

$$\sinh ix = \frac{1}{2}(e^{+ix} - e^{-ix})$$

$$\tanh ix = \frac{(e^{+ix} - e^{-ix})}{(e^{+ix} + e^{-ix})}$$

$$\cosh ix = \cos x$$

$$\sinh ix = i \sin x$$

$$\tanh ix = i \tan x$$

With imaginary arguments

$$\cos x = \frac{1}{2}(e^{+ix} + e^{-ix})$$



$$\sin x = \frac{1}{2i}(e^{+ix} - e^{-ix})$$



$$\tan x = \frac{1}{i} \frac{(e^{+ix} - e^{-ix})}{(e^{+ix} + e^{-ix})}$$



$$\cosh x = \frac{1}{2}(e^{+x} + e^{-x})$$

$$\sinh x = \frac{1}{2}(e^{+x} - e^{-x})$$

$$\tanh x = \frac{(e^{+x} - e^{-x})}{(e^{+x} + e^{-x})}$$

X

$$\cos ix = \cosh x$$

$$\sin ix = i \sinh x$$

$$\tan ix = i \tanh x$$

$$\cosh ix = \cos x$$

$$\sinh ix = i \sin x$$

$$\tanh ix = i \tan x$$

i X

Euler Formula

Euler Formula

$$e^{+ix} = \cos x + i \sin x$$

$$e^{-ix} = \cos x - i \sin x$$

Euler Formula

$$e^{+ix} = \cosh ix + \sinh ix$$

$$e^{-ix} = \cosh ix - \sinh ix$$

$$\cos ix = \cosh x$$

$$\sin ix = i \sinh x$$

$$\tan ix = i \tanh x$$

$$\cosh ix = \cos x$$

$$\sinh ix = i \sin x$$

$$\tanh ix = i \tan x$$

Modulus of $\sin(z) - 1$

$$\begin{aligned}\sin(z) &= \sin(x + iy) \\ &= \sin(x)\cos(iy) + \cos(x)\sin(iy) \\ &= \sin(x)\cosh(y) + i\cos(x)\sinh(y)\end{aligned}$$

$$\begin{aligned}|\sin(z)|^2 &= \sin(z) \overline{\sin(z)} \\ &= \frac{1}{2i}(e^{+i(x+iy)} - e^{-i(x+iy)}) \frac{-1}{2i}(e^{-i(x-iy)} - e^{+i(x-iy)}) \\ &= \frac{1}{4}(e^{-y+ix} - e^{+y-ix})(e^{-y-ix} - e^{+y+ix}) \\ &= \frac{1}{4}(e^{-2y} - e^{+2ix} - e^{-2ix} + e^{+2y}) \\ &= \frac{1}{4}(e^{+2y} - 2 + e^{-2y} - e^{+2ix} + 2 - e^{-2ix}) \\ &= +\frac{1}{4}(e^{+2y} - 2 + e^{-2y}) - \frac{1}{4}(e^{+2ix} - 2 + e^{-2ix}) \\ &= \left[\frac{1}{2}(e^{+y} - e^{-y})\right]^2 + \left[\frac{1}{2i}(e^{+ix} - e^{-ix})\right]^2 \\ &= \sin^2(x) + \sinh^2(y)\end{aligned}$$

Modulus of $\sin(z)$ - (2)



$$\begin{aligned}\sin(z) &= \sin(x + iy) \\ &= \sin(x)\cos(iy) + \cos(x)\sin(iy) \\ &= \sin(x)\cosh(y) + i\cos(x)\sinh(y)\end{aligned}$$

$$\begin{aligned}|\sin(z)|^2 &= |\sin(x)\cosh(y) + i\cos(x)\sinh(y)|^2 \\ &= \sin^2(x)\cosh^2(y) + \cos^2(x)\sinh^2(y) \\ &= \sin^2(x)(1 + \sinh^2(y)) + (1 - \sin^2(x))\sinh^2(y) \\ &= \sin^2(x) + \sin^2(x)\sinh^2(y) + \sinh^2(y) - \sin^2(x)\sinh^2(y) \\ &= \sin^2(x) + \sinh^2(y)\end{aligned}$$

$$\cosh^2 \alpha - \sinh^2 \alpha = 1$$

$$\cos^2 \alpha + \sin^2 \alpha = 1$$

Graphs of $\sin(z)$



$$\sin(z) = \sin(x+iy)$$


$$= \sin(x) \cosh(y) + i \cos(x) \sinh(y)$$

$$|\sin(z)|^2 = \sin^2(x) + \sinh^2(y)$$

<http://en.wikipedia.org/>

Domain Coloring – Argument

Domain Coloring



Argument of $f(z)$

hue to phase/angle/argument
legend:


hue	phase (radians)
red	$0 \text{ mod } 2\pi$
yellow	$\pi/3 \text{ mod } 2\pi$
green	$2\pi/3 \text{ mod } 2\pi$
cyan	$\pi \text{ mod } 2\pi$
blue	$4\pi/3 \text{ mod } 2\pi$
magenta	$5\pi/3 \text{ mod } 2\pi$

$\theta = \pi \text{ mod } 2\pi$ —

Each discontinuity
in intensity occurs
when $|z|=2^n$, for
integer $n (0, -1, -2, \dots)$

The Unit Circle



$\theta = \pi/2 \text{ mod } 2\pi$



— $\theta = 0 \text{ mod } 2\pi$



<http://en.wikipedia.org/>

Domain Coloring – Modulus




<http://en.wikipedia.org/>

Domain Coloring of $\sin(z)$




Domain coloring of $\sin(z)$ over $(-\pi, \pi)$ on x and y axes. Brightness indicates absolute magnitude, saturation represents imaginary and real magnitude.




<http://en.wikipedia.org/>

Another domain coloring of $\sin(z)$



<http://en.wikipedia.org/>

Domain Coloring of $\sin(x)$, $\cos(x)$, $\tan(x)$



$$\sin(z) = \sin(x+iy)$$

$$\cos(z) = \cos(x+iy)$$

$$\tan(z) = \tan(x+iy)$$

<http://en.wikipedia.org/>

References

- [1] <http://en.wikipedia.org/>
- [2] J. S. Walther, A Unified Algorithm for Elementary Functions
- [3] J. Calvert, <http://mysite.du.edu/~jcalvert/math/hyperb.htm>